

Recurrent Proofs of the Irrationality of Certain Trigonometric Values

Li Zhou and Lubomir Markov

In this note we exploit recurrences of integrals to give new elementary proofs of the irrationality of $\tan r$ for $r \in \mathbb{Q} \setminus \{0\}$ and $\cos r$ for $r^2 \in \mathbb{Q} \setminus \{0\}$. We also discuss applications of our technique to simpler irrationality proofs such as those for π , π^2 , and certain values of exponential and hyperbolic functions.

1 Irrationality of $\tan r$ for $r \in \mathbb{Q} \setminus \{0\}$.

For a nonzero rational r , the irrationality of $\tan r$ was first proved by J. H. Lambert in 1761 by means of continued fractions [1, pp. 129–146]. We now present a new direct proof using a recurrence for an integral.

Theorem 1. *$\tan r$ is irrational for nonzero rational r .*

Proof. The irrationality of π will be a by-product of this proof, so we start by supposing that $r \in \mathbb{Q} \setminus \{k\pi : k \in \mathbb{Z}\}$. Write $r = a/b$ with $a, b \in \mathbb{Z}$ and assume that $\tan(r/2) = p/q$ with $p, q \in \mathbb{Z}$. For $n \geq 0$, let $f_n(x) = (rx - x^2)^n/n!$ and $I_n = \int_0^r f_n(x) \sin x \, dx$. Then $b^n I_n \rightarrow 0$ as $n \rightarrow \infty$, $I_0 = 1 - \cos r$, and $I_1 = 2(1 - \cos r) - r \sin r$. Integrating by parts twice, we get that for $n \geq 2$,

$$I_n = - \int_0^r f_n''(x) \sin x \, dx = (4n - 2)I_{n-1} - r^2 I_{n-2}. \quad (1)$$

By inducting on n using (1), we see that for $n \geq 0$, $I_n = u_n(1 - \cos r) + v_n \sin r$, where u_n and v_n are polynomials in r with integer coefficients and degrees at most n . Moreover, if two consecutive terms of the sequence $\langle I_n \rangle$ are 0, then (1) forces all terms of $\langle I_n \rangle$ to be 0, and in particular $I_0 = 0$, a contradiction. Hence $\langle I_n \rangle$ has infinitely many nonzero terms. Therefore, we can pick a large enough n so that $b^n q \csc r I_n = b^n q [u_n \tan(r/2) + v_n]$ is a nonzero integer in $(-1, 1)$, a contradiction.

Notice that $\tan(\pi/4) = 1$, so $\pi/2 \notin \mathbb{Q}$, which implies that $\mathbb{Q} \setminus \{k\pi : k \in \mathbb{Z}\} = \mathbb{Q} \setminus \{0\}$. Thus we have proved that $\tan(r/2) \notin \mathbb{Q}$ for all $r \in \mathbb{Q} \setminus \{0\}$. \square

A closer inspection of our proof reveals that u_n and v_n/r are polynomials in r^2 . So a slightly stronger conclusion can be squeezed out of the proof, namely that $(\tan r)/r$ is irrational whenever $r^2 \in \mathbb{Q} \setminus \{0\}$. This stronger result was first established through a different elementary approach by Inkeri [2].

2 Simpler Proofs of Irrationality.

From the previous proof we see that recurrence is a double-edged sword. It is sharp and swift in showing that a sequence is integer-valued and has an infinite nonzero subsequence. We can also use recurrence to give similar proofs of the irrationality of π , π^2 , e^r , etc.. However, in these easier cases, the corresponding sequences of integrals are positive, so there is no need to argue for the existence of a nonzero subsequence. Consequently the proofs can be really short and charming. For a flavor of it, we present a direct proof of the irrationality of π which is even shorter than the celebrated one-page proof given by Niven [3].

Theorem 2. π is irrational.

Proof. Assume that $\pi = a/b$ with $a, b \in \mathbb{N}$. Let $f_n(x) = (\pi x - x^2)^n/n!$ and $I_n = \int_0^\pi f_n(x) \sin x dx$. Then $b^n I_n \rightarrow 0$ as $n \rightarrow \infty$, $I_0 = 2$, and $I_1 = 4$. Replacing r by π in (1) we get $I_n = (4n - 2)I_{n-1} - \pi^2 I_{n-2}$ for $n \geq 2$. By induction on n using this recurrence, we see that for $n \geq 0$, I_n is a polynomial in π with integer coefficients and degree at most n . Hence for a large enough n , $b^n I_n$ is an integer in $(0, 1)$, a contradiction. \square

Notice that the terms of $\langle I_n \rangle$ are really polynomials in π^2 , so our proof only needs very minor changes to show the stronger conclusion that π^2 is irrational. In fact in [5], Schröder presented a very similar proof by recurrence of the irrationality of π^2 .

For the irrationality of e^r for nonzero rational r , the interested reader can imitate the process with the sequence $I_n = \int_0^r f_n(x) e^x dx$, where $f_n(x) = (rx - x^2)^n/n!$ as well.

3 Irrationality of $\cos r$ for $r^2 \in \mathbb{Q} \setminus \{0\}$.

Next we turn our attention to the cosine function. The classical elementary proof of the irrationality of $\cos r$ for nonzero rational r was given by Niven [4, Theorem 2.5, pp. 16–19]. Niven's proof can be slightly modified to show that $\cos r$ is irrational whenever $r^2 \in \mathbb{Q} \setminus \{0\}$, as observed by Inkeri [2]. We now use the full force of recurrence to give a proof which is more direct than Inkeri's modification.

Theorem 3. If $r^2 \in \mathbb{Q} \setminus \{0\}$ then $\cos r$ is irrational.

Proof. Assume that $r^2 = a/b$ with $a, b \in \mathbb{Z} \setminus \{0\}$ and $\cos r = p/q$ with $p, q \in \mathbb{Z}$. For $n \geq 0$, let $f_n(z) = (r^2 z^2 - z^4)^n/n!$, $I_n = \int_0^r f_n(z) \sin(r - z) dz$, $J_n = \int_0^r z f_n(z) \cos(r - z) dz$, $K_n = \int_0^r z^2 f_n(z) \sin(r - z) dz$, and $L_n = \int_0^r z^3 f_n(z) \cos(r - z) dz$. Then b^{2n+1} times each of the four integrals approaches 0 as $n \rightarrow \infty$. Direct integration yields $I_0 = 1 - \cos r = J_0$, $K_0 = r^2 - 2 + 2 \cos r$, and $L_0 = 3K_0$.

Integrating each integral by parts once, we get that for $n \geq 1$,

$$I_n = 4L_{n-1} - 2r^2 J_{n-1}, \quad (2)$$

$$J_n = (4n+1)I_n - 2r^2 K_{n-1}, \quad (3)$$

$$K_n = -(4n+2)J_n + 2r^2 L_{n-1}, \quad (4)$$

$$L_n = (4n+3)K_n + 2nr^2 I_n - 2r^4 K_{n-1}. \quad (5)$$

Induction on n in these recurrences implies that for $n \geq 0$, the four sequences have the form $u_n + v_n \cos r$, where u_n and v_n are polynomials in r^2 with integer coefficients and degrees at most $2n+1$. Moreover, suppose that $I_m = J_m = K_m = L_m = 0$ for some $m \geq 1$. Then (3) and (4) imply that $K_{m-1} = 0$ and $L_{m-1} = 0$. Thus (2) yields $J_{m-1} = 0$. Eliminating K_{n-1} from (3) and (5), we see that for $n \geq 1$, I_n can be expressed in terms of J_n , K_n , and L_n ; and keep in mind also that $I_0 = J_0$. Hence $I_{m-1} = 0$. By this argument of infinite descent we conclude that $I_0 = J_0 = K_0 = L_0 = 0$, which contradicts the fact that $2I_0 + K_0 = r^2 \neq 0$. Therefore, at least one of the four sequences has infinitely many nonzero terms. Pick such a sequence and a large enough n so that $b^{2n+1}q$ times the corresponding integral is a nonzero number in $(-1, 1)$, while it also has the form $b^{2n+1}q(u_n + v_n \cos r)$ which is an integer, a contradiction. Thus $\cos r \notin \mathbb{Q}$ whenever $r^2 \in \mathbb{Q} \setminus \{0\}$. \square

Corollary 1. π^2 is irrational. Also, if $r^2 \in \mathbb{Q} \setminus \{0\}$ then $\sin^2 r$, $\cos^2 r$, and $\tan^2 r$ are all irrational.

Proof. The claims follow immediately from Theorem 3 and the identities $\cos \pi = -1$ and $\cos 2r = 1 - 2\sin^2 r = 2\cos^2 r - 1 = (1 - \tan^2 r)/(1 + \tan^2 r)$. \square

Finally, the observant reader perhaps has noticed that our proof of Theorem 3 allows r^2 to be a negative rational number. Since $\cosh r = \cos(ir)$, all the analogous statements about hyperbolic functions are included in our results. The skeptical reader is invited to work out the details by substituting $r = it$ and $z = iy$ into I_n , J_n , K_n , and L_n . In fact, the resulting real integrals are nonzero and thus the proof is shorter, because the argument of infinite descent is not needed.

Acknowledgments. We are grateful to our friend Prof. Paul Yiu whose comments on an earlier draft led to a substantial improvement of the paper. We would also like to thank the anonymous referees for their suggestions and for informing us of the references [2] and [5].

References

- [1] L. Berggren, J. Borwein, and P. Borwein, eds., *Pi: A Source Book*, 3rd ed., Springer, New York, 2004.
- [2] K. Inkeri, Eräiden transsendenttifunktioiden arvojen irrationaalisuudesta, *Arkhimedes*, no. 1 (1965) 15–21.

- [3] I. Niven, A simple proof that π is irrational, *Bull. Amer. Math. Soc.* **53** (1947) 509.
- [4] ———, *Irrational Numbers*, Mathematical Association of America, Washington, DC, 1956.
- [5] E. M. Schröder, Zur irrationalität von π^2 und π , *Mitt. Math. Ges. Hamburg* **13** (1993) 249.

Department of Mathematics, Polk State College, Winter Haven, FL 33881
lzhou@polk.edu

Department of Mathematics and CS, Barry University, Miami Shores, FL 33161
lmarkov@mail.barry.edu